

Morse Theory on Triangulated Meshes

MUSTAFA HAJIJ



What is Morse theory ?

Morse Theory is a tool in differential topology that is concerned with the relations between the geometric and topological aspects of manifolds and the real-valued functions defined on them.


What is Morse theory ?

Morse Theory is a tool in differential topology that is concerned with the relations between the geometric and topological aspects of manifolds and the real-valued functions defined on them.

One of the primary interests in this theory is the relationship between the topology of a smooth manifold M and the critical points of a real-valued smooth function f defined on M .

What is Morse theory ?


Consider the following example. Let M be a 2-dimensional torus.



What is Morse theory ?


Consider the following example. Let M be a 2-dimensional torus.

Define a function f on M to be the function that sends every point (x,y,z) on the torus to z coordinate.



What is Morse theory ?


Consider the inverse image of the value z_1 under the function f .



What is Morse theory ?

Consider the inverse image of the value z_1 under the function f .

As we increase z_1 to z_2 , the topology of the level set changes from one circle to 2 circles




What is Morse theory ?

Consider the inverse image of the value z_1 under the function f .


As we increase z_1 to z_2 , the topology of the level set changes from one circle to 2 circles

We call the point on M at which this topological change occurs a *critical point*



What is Morse theory ?

Morse theory studies carefully the topological changes that happen to the manifold M as we pass from a critical point.



Morse theory on smooth surfaces

Let M be a compact smooth surface and let $I = [a, b] \subseteq \mathbb{R}$, where $a < b$, be a closed interval. Let $f : M \rightarrow I$ be a smooth function defined on M . A point $x \in M$ is called a *critical point* of f if the differential df_x is zero. A value c in \mathbb{R} is called a *critical value* of f if $f^{-1}(c)$ contains a critical point of f . A point in M is called a *regular point* if it is not a critical point. Similarly, if a value $c \in \mathbb{R}$ is not a critical value then we call it a regular value. The inverse function theorem implies that for every regular value c in \mathbb{R} the level set $f^{-1}(c)$ is a 1-manifold, i.e., $f^{-1}(c)$ is a disjoint union of circles. A critical point is called *non-degenerate* if the matrix of the second partial derivatives of f , called the *Hessian matrix*, is non-singular. If all the critical points of f are non-degenerate and all critical points have distinct values then f is a *Morse function*.


Morse Lemma

Lemma (Morse Lemma) Let M be a smooth surface, $f : M \rightarrow \mathbb{R}$ be a smooth function and p be a non-degenerate critical point of f . We can choose a chart (ϕ, U) around p such that $f \circ \phi^{-1}$ takes exactly one of the following three forms:

1. $f \circ \phi^{-1}(X, Y) = X^2 + Y^2 + c.$

2. $f \circ \phi^{-1}(X, Y) = -X^2 - Y^2 + c.$


3. $f \circ \phi^{-1}(X, Y) = X^2 - Y^2 + c.$



minimum



saddle




maximum

The form of a Morse function f around a critical point can be proven to be independent of the choice of the chart.

Handle decomposition

Let M be a smooth surface and let $f : M \rightarrow \mathbb{R}$ be a Morse function defined on M . Define the set

$$M_{f,t} = \{x \in M : f(x) \leq t\}.$$




Handle decomposition

Theorem *Let $f : M \rightarrow \mathbb{R}$ be Morse function. Let p be a critical point of index i and $f(p) = t$ be its corresponding critical value. Let ϵ be chosen small enough so that f has no critical values on the interval $[t - \epsilon, t + \epsilon]$. Then :*

- 1. If $\text{index}_f(p) = 0$, then $M_{t+\epsilon}$ is diffeomorphic to the disjoint union of $M_{t-\epsilon}$ and a 2-dimensional disk D^2 .*
- 2. If $\text{index}_f(p) = 1$, then $M_{t+\epsilon}$ can be obtained from $M_{t-\epsilon}$ by attaching a 1-handle. That means that $M_{t+\epsilon}$ can be obtained by gluing a rectangular strip $D^1 \times D^1$ to the boundary of $M_{t-\epsilon}$ along $D^1 \times \partial D^1$.*
- 3. If $\text{index}_f(p) = 2$, then $M_{t+\epsilon}$ can be obtained by capping off the surface $M_{t-\epsilon}$ with a disk D^2 . That means that $M_{t+\epsilon}$ is obtained by gluing a disk D^2 along its boundary ∂D^2 to one of the boundary components of $M_{t-\epsilon}$.*

Morse Theory on Meshes


In practice we approximate surfaces by triangulated meshes.




Examples of triangulated meshes

Morse Theory on Meshes

In practice we approximate surfaces by triangulated meshes.



Mesh components




Examples of triangulated meshes

Morse Theory on Meshes

Suppose that we have a mesh M and suppose that f is a scalar function defined on the set of vertices of M .

Morse Theory on Meshes


Suppose that we have a mesh M and suppose that f is a scalar function defined on the set of vertices of M .




Consider a local ring neighborhood around a vertex and consider the values of v on the ring

Morse theory on Meshes

Suppose that we have a mesh M and suppose that f is a scalar function defined on the set of vertices of M .




Consider a local ring neighborhood around a vertex and consider the values of v on the ring




Locally, the Morse scalar function around a vertex has one of the following possibilities


Morse Scalar Functions on Meshes




regular



maximum




minimum




saddle


Morse Scalar Functions on Meshes




regular




maximum




minimum




saddle



minimum



maximum



saddle

Types of vertices

Given a Morse function f on a triangulated mesh M .
Then we can classify the vertices of M as follows

The upper link of v is defined as


$$Lk^+(v) = \{u \in Lk(v) : f(u) > f(v)\},$$

and the lower link is defined by

$$Lk^-(v) = \{u \in Lk(v) : f(u) < f(v)\},$$

and mixed link

$$Lk^\pm(v) = \{(u_1, u_2) : f(u_1) < f(v) < f(u_2)\}.$$









v is *regular* if $|Lk^\pm(v)| = 2$

is a *maximum* with index 2 if $|Lk^+(v)| = 0$,

is a *minimum* with index 0 if $|Lk^-(v)| = 0$

is a *saddle* with index 1 and multiplicity $m \geq 1$ if $|Lk^\pm(v)| = 2 + 2m$

Definition: A scalar function f on a triangulated mesh M is PL Morse function if each vertex is either regular or simple critical (minimum, maximum or saddle with $m=1$) and the function values of the vertices are distinct.



Morse Scalar Functions on Meshes

We will represent the isolines as follows:





Morse Scalar Functions on Meshes

In the following example we represent :

minimum point by a blue sphere
maximum point by a red sphere
saddle point by a green sphere



Morse Scalar Functions on Meshes




Applications of Morse Theory

Morse theory has found many applications recently in geometric processing

Applications of Morse Theory

Morse theory has found many applications recently in geometric processing


Extracting Surface feature lines (Sahner et. al.)




Applications of Morse Theory

Morse theory has found many applications recently in geometric processing

Extracting Surface feature lines (Sahner et. al.)




Surface quadrangulation (Dong et. al.)




Applications of Morse Theory

Morse theory has found many applications recently in geometric processing

Extracting Surface feature lines (Sahner et. al.)




Surface quadrangulation (Dong et. al.)



Cutting a surface into a disk (Ni et. al.)

Reeb Graph

Given a surface M and a scalar function defined on it, we can define a combinatorial structure called the *Reeb graph* of M and f by collapsing the level sets of f as illustrated in the figure



Thank You